

ADVANCED TOPICS IN ALGEBRAIC TOPOLOGY — EXERCISE SHEET 8

Exercise class: Friday, 19th of June, 11-12

Website with further material, including exercise sheets:

<https://www.groups.ma.tum.de/algebra/scheimbauer/advanced-topics-in-algebraic-topology/>

- Exercise 1.** (a) Use the Künneth formula to compute the cohomology of $T^n := \overbrace{S^1 \times \cdots \times S^1}^{n \text{ times}}$.
Give an explicit basis!
- (b) What is the Euler characteristic (as defined on the last sheet) of T^n ?
- (c) Determine the Poincaré dual $\eta_S \in H^1(T^2)$ of the submanifold $S := S^1 \times \{1\} \subset T^2$.

Definition 1 (Real and complex projective lines). The **complex projective line** is the topological space

$$\mathbb{C}\mathbb{P}^1 := \frac{\mathbb{C}^2 \setminus \{0\}}{\sim}$$

(with the quotient topology), where $v \sim w$ if and only if v and w span the same vector subspace $\langle v \rangle_{\mathbb{C}} = \langle w \rangle_{\mathbb{C}}$ of \mathbb{C}^2 .

Replacing \mathbb{C} by \mathbb{R} yields the **real projective line** $\mathbb{R}\mathbb{P}^1$.

- Exercise 2.** (a) Equip $\mathbb{R}\mathbb{P}^1$ with the structure of a smooth manifold diffeomorphic to S^1 .
- (b) Equip $\mathbb{C}\mathbb{P}^1$ with the structure of a smooth manifold which is diffeomorphic to the 2-sphere S^2 .
- Exercise 3** (Hopf fibration). (a) Show that the projection $p: \mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{C}\mathbb{P}^1$ is a non-trivial fiber bundle with fibers diffeomorphic to \mathbb{C}^\times .
- (b) Show that p contains a non-trivial fiber subbundle $S^3 \rightarrow S^2 \cong \mathbb{C}\mathbb{P}^1$ with fibers diffeomorphic to S^1 .

- Exercise 4** (Tautological bundle on projective lines). (1) Construct a non-trivial real vector bundle of rank 1 on $\mathbb{R}\mathbb{P}^1$ such that the fiber over each point $[v] \in \mathbb{R}\mathbb{P}^1$ is the \mathbb{R} -vector space $\langle v \rangle_{\mathbb{R}} \subset \mathbb{R}^2$. Have you seen this bundle before?
- (2) Construct a non-trivial complex vector bundle of rank 1 on $\mathbb{C}\mathbb{P}^1$ such that the fiber over each point $[v] \in \mathbb{C}\mathbb{P}^1$ is the \mathbb{C} -vector space $\langle v \rangle_{\mathbb{C}} \subset \mathbb{C}^2$. How does this bundle relate to the one constructed in Exercise 3?

Exercise 5 (Bonus challenge). Let M be a compact non-orientable manifold of dimension n . Prove that $H^n(M) = 0$. (Hint: Consider the orientation double cover from sheet 5.)